

MICROWAVE EXTRACTION AND  
MICROENCAPSULATION OF POLYPHENOL  
FROM *PHYLLANTHUS NIRURI*

NGUANG SUOK LING

MASTER OF SCIENCE

UNIVERSITI MALAYSIA PAHANG



## **SUPERVISOR'S DECLARATION**

I hereby declare that I have checked this thesis and in my opinion, this thesis is adequate in terms of scope and quality for the award of the degree of Master of Science in Chemical Engineering.

---

(Supervisor's Signature)

Full Name : DR. JOLIUS BIN GIMBUN

Position : ASSOCIATE PROFESSOR

Date :



## **STUDENT'S DECLARATION**

I hereby declare that the work in this thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

---

(Student's Signature)

Full Name : NGUANG SUOK LING

ID Number : MKC16003

Date :

MICROWAVE EXTRACTION AND MICROENCAPSULATION OF  
POLYPHENOL FROM *PHYLLANTHUS NIRURI*

NGUANG SUOK LING

Thesis submitted in fulfillment of the requirements  
for the award of the degree of  
Master of Science/Master of Engineering

Faculty of Chemical & Natural Resources Engineering  
UNIVERSITI MALAYSIA PAHANG

NOVEMBER 2018

## **ACKNOWLEDGEMENTS**

I would like to extend my gratitude and appreciation to the following people and organizations. My supervisor, Assoc. Prof. Dr. Jolius Gimbun for the guidance and encouragement, sage advice, numerous comments which make this thesis reality. I appreciate your expertise and enthusiasm in all the experiments that I had carried out throughout this study. I would like to extend my appreciation towards Pang Sook Fun who acts as my tutor, who guided me all the way for this research study and solved all the difficulties I faced in the lab. Apart from that, I would like to thanks to Yeong Yi Ling, and Kee Kieng Lee, who assist me throughout the lab experiment activities and also contributed 6 awards and 3 papers.

## ABSTRAK

Dukung anak, mengandungi pelbagai komponen bioaktif yang menyumbang dalam bidang perubatan. Ia mempunyai sifat antioksidan, anti radang, anti kancer dan dapat mngubati Hepatitis B dan jangkitan usus penyakit kuning. Pengekstrakan ialah kaedah yang biasa digunakan untuk mendapatkan komponen bioaktif daripada bahan-bahan tumbuhan. Hasil komponen bioaktif itu bergantung kepada kaedah pengekstrakan, pelarut dan keadaan pengekstrakan. Dalam karya ini, kaedah pengekstrakan ultrasonik dan kaedah pengekstrakan gelombang mikro dikaji kerana kekurangan penyelidikan sebelum ini mengenai pengekstrakan ultrasonik dan pengekstrakan gelombang mikro telah disediakan dalam kesusasteraan. Dalam pendapatan kajian ini, didapati bahawa hasil komponen bioaktif adalah sangat bergantung kepada kekutuban pelarut yang digunakan dalam pengekstrakan, hasil tertinggi phyllanthin (4.56mg Phy/g DW) telah diperolehi dengan menggunakan 20% akueus Isopropanol manakala hasil tertinggi quercetin (10.14mg Que / g DW) telah diperolehi dengan menggunakan 20% etanol berair dan hasil tertinggi asid Gallic (15.44mg GAE / g DW) telah diperolehi dengan menggunakan air. Kekutuban pelarut meningkatkan pengeluaran kedua-dua komponen hydroxylated dan methoxylated dari *P. niruri*. Daripada analisis reka bentuk komposit pusat, pengekstrakan gelombang mikro pada kuasa pengeluaran di 250W, masa pengekstrakan dalam 2.47 minit hingga 5.72 minit dan kepekatan etanol daripada 36.58% hingga 76.31% mampu untuk mendapatkan hasil optimum pengekstrakan polifenol dengan kebaikan 91.70%. Pengekstrakan gelombang mikro disediakan pengekstrakan yang cepat tanpa ketara menjejaskan hasil pengekstrakan. produk tepung sering dikehendaki kerana jangka hayat yang lebih panjang, mudah untuk penggunaan dan mudah pengangkutan / pengendalian. Proses membuat serbuk sering dijalankan di dalam pengering semburan pada suhu tinggi (180° C). Untuk mengurangkan degradasi komponen bioaktif semasa pengeringan semburan, kaedah pemikrokapsulan diperkenalkan. WPI dan MD serbuk semburan encapsulation telah mengadakan pengekal polifenol yang baik dari *P. niruri*. Pemikrokapsulan menggunakan campuran WPI dan MD pada nisbah 1: 9 menyumbang pengekal tertinggi phyllanthin (84,33%), asid Gallic (88,93%) dan quercetin (88.39%) diikuti oleh MD dan WPI pengkapsulan. Pemikrokapsulan menggunakan campuran WPI dan MD pada nisbah 1: 9 dicadangkan kerana ia menyediakan pemeliharaan yang lebih baik daripada polifenol semasa pengeringan semburan yang bertentangan dengan merangkumi protein tunggal WPI dan MD. Keputusan analisis oleh UPLC menggambarkan asid Gallic dan quercetin adalah lebih mudah terdedah kepada pencemaran haba daripada phyllanthin semasa pengeringan semburan. Kajian yang lebih mendalam terhadap ujian toxic dicadangkan dengan menggunakan binatang untuk memastikan *P. niruri* ekstrak tidak membahayakan dalam ACCU lab yang diiktirafkan.

## ABSTRACT

*Phyllanthus Niruri* (ver. name: Dukung Anak) contains miscellaneous bioactive compounds which contribute in various medical effects such as antioxidant, anti-inflammatory, anti-cancer and treating Hepatitis-B, jaundice intestinal infection. Extraction is the most common method to obtain the bioactive component from the plant materials. The yield of bioactive component in the extract is dependent on the solvent used, extraction method and condition. In this work, ultrasonic assisted extraction and microwave assisted extraction method were studied as there's limited work on ultrasonic assisted extraction and microwave assisted extraction were available in the literature. From the finding of this work, it is found that the yield of the bioactive component is highly dependent on solvent polarity used in the extraction, the highest yield of phyllanthin (4.56mg Phy/g DW) was obtained using 20% aqueous Isopropanol whereas the highest yield of quercetin (10.14mg Que/g DW) was obtained by using 20% aqueous ethanol and highest yield of gallic acid (15.44mg GAE/g DW) was obtained by using water. The polarity of solvent enhances the extraction of both hydroxylated and methoxylated compounds from the *P. niruri*. From the central composite design analysis, microwave assisted extraction at extraction power at 250W, extraction time ranged from 2.47 minutes to 5.72 minutes and ethanol concentration will ranged from 36.58% to 76.31% able to obtain the optimum yield of polyphenol extraction with the desirability of 91.70%. Microwave assisted extraction provided a fast extraction without significantly compromising the extraction yield. Powdered product is often desired due to its longer lifespan, convenient for consumption and easier transportation/ handling. The process of powder making is often carried out in a spray dryer at high temperature (180°C). To minimize degradation of the bioactive compounds during spray drying, microencapsulation method was introduced. WPI and MD encapsulation spray powder had performed a good polyphenol retention from *P. niruri*. Microencapsulation using mixture of WPI and MD at the ratio 1:9 delivered highest retention of phyllanthin (84.33%), gallic acid (88.93%) and quercetin (88.39%) followed by MD and WPI encapsulation. Microencapsulation using mixture of WPI and MD at ratio 1:9 is suggested as it provides a better preservation of polyphenol during spray drying as opposed to single protein encapsulate of WPI and MD. The results analysis by UPLC illustrates gallic acid and quercetin are more susceptible to thermal degradation than phyllanthin during spray drying. It is recommended to perform a toxicity analysis of *P. niruri* extract in ACCU accredited lab before it can undergo clinical trial.

## **TABLE OF CONTENT**

**DECLARATION**

**TITLE PAGE**

**ACKNOWLEDGEMENTS** **ii**

**ABSTRAK** **iii**

**ABSTRACT** **iv**

**TABLE OF CONTENT** **v**

**LIST OF TABLES** **ix**

**LIST OF FIGURES** **xi**

**LIST OF SYMBOLS** **xiii**

**LIST OF ABBREVIATIONS** **xiv**

**CHAPTER 1 INTRODUCTION** **1**

1.1 Background 1

1.2 Problem statement 2

1.3 Objectives 5

1.4 Scope of this research 5

1.5 Main Contribution of this work 6

1.6 Summary 6

**CHAPTER 2 LITERATURE REVIEW** **7**

2.1 Overview 7

2.2 Introduction 7

2.3 Extraction method for flavonoids and phenolic content in plant 8



2.3.1	Ultrasonic assisted extraction	9
2.3.2	Microwave assisted extraction	11
2.4	Bioactive compounds from <i>phyllanthus niruri</i>	14
2.5	Spray Drying	16
2.6	Microencapsulation	19
2.7	Ultra Performance Liquid Chromatography	20
2.8	Summary	23
<b>CHAPTER 3 METHODOLOGY</b>		<b>24</b>
3.1	Overview	24
3.2	Chemicals	25
3.3	Plant Material	25
3.4	Extraction Method	26
3.4.1	Ultrasonic Assisted Extraction (UAE)	26
3.4.2	Microwave Assisted Extraction (MAE)	28
3.5	Microencapsulation and Spray Drying	31
3.5.1	Encapsulating agent solution preparation	31
3.5.2	Spray Drying Process	32
3.6	Analysis	33
3.6.1	Total Phenolic Content	33
3.6.2	Total Flavonoids Content	34
3.6.3	Antioxidant Activity	34
3.6.4	Ultra performance liquid chromatography	35
3.6.5	Scanning electron microscopy (SEM)	37
3.6.6	Particle size analysis	38
3.6.7	Moisture content analysis	38

3.7	Statistical Analysis	38
<b>CHAPTER 4 RESULTS AND DISCUSSION</b>		<b>39</b>
4.1	Overview	39
4.2	Introduction	39
4.3	UPLC Quantification of polyphenol	40
4.4	Influence of solvent type to the polyphenols extraction	42
4.5	Factorial Analysis on Ultrasonic Assisted Extraction (UAE)	44
4.5.1	Effect of Solvent purity, Time and Amplitude on UAE	44
4.5.2	Optimization on the polyphenol extraction	47
4.6	Factorial Analysis on Microwave Assisted Extraction (MAE)	55
4.6.1	Effect of Time, Power, Solvent purity and Solid Liquid Ratio on MAE.	55
4.6.2	Optimizing on polyphenol extraction by MAE	61
4.7	Summary	68
<b>CHAPTER 5 MICROENCAPSULATION</b>		<b>70</b>
5.1	Introduction	70
5.2	Moisture content	70
5.3	Particle Size Distribution	71
5.4	Particle morphology	72
5.5	Microencapsulation of polyphenolic compounds from <i>P. niruri</i>	74
5.6	Summary	75
<b>CHAPTER 6 CONCLUSION AND RECOMMENDATION</b>		<b>76</b>
6.1	Conclusion	76
6.2	Recommendation	77

<b>REFERENCES</b>	<b>79</b>
<b>APPENDIX A</b>	<b>88</b>

## LIST OF TABLES

Table 2.1	Previous work on <i>Phyllanthus Niruri</i> .	8
Table 2.2	Physical constants and dissipation factors for solvents usually used in microwave-assisted extraction (MAE) ( Zlotorzynski 1995 and Jassie et al., 1997).	12
Table 2.3	Review of spray drying condition.	18
Table 2.4	Review of HPLC analysis for <i>Phyllanthus Niruri</i> .	21
Table 3.1	Factors and their designated low and high value.	27
Table 3.2	Experimental design matrix for UAE.	27
Table 3.3	Levels of extraction condition variables tested in CCD.	28
Table 3.4	Experimental design matrix for optimization.	28
Table 3.5	Factors and their designated low and high value.	29
Table 3.6	Experimental design matrix for MAE.	29
Table 3.7	Levels of extraction condition variables tested in CCD.	30
Table 3.8	Experimental design matrix for optimization.	31
Table 3.9	Encapsulating agent and its concentration.	32
Table 4.1	Effect of solvent on polyphenols extraction from <i>Phyllanthus Niruri</i> .	43
Table 4.2	Experimental design and response for factorial analysis of UAE.	44
Table 4.3	Sum of squares and the percent contribution for each term for Phyllanthin.	45
Table 4.4	Sum of squares and the percent contribution for each term for Gallic acid.	45
Table 4.5	Sum of squares and the percent contribution for each term for Quercetin.	46
Table 4.6	Sum of squares and the percent contribution for each term for Total Phenolic Content.	46
Table 4.7	Sum of squares and the percent contribution for each term for Total Flavonoid Content.	46
Table 4.8	Sum of squares and the percent contribution for each term for Antioxidant Activity.	47
Table 4.9	Suggested best condition for factors in UAE for maximizing all responses.	47
Table 4.10	Experimental design and response for optimization.	49
Table 4.11	Condition for factors optimizing polyphenol extraction.	53
Table 4.12	Comparison between predicted and experimental value for optimum condition.	54
Table 4.13	Experimental design and response for MAE factorial analysis.	56

Table 4.14 Sum of squares and the percent contribution for each term for phyllanthin	58
Table 4.15 Sum of squares and the percent contribution for each term for gallic acid.	58
Table 4.16 Sum of squares and the percent contribution for each term for quercetin.	58
Table 4.17 Sum of squares and the percent contribution for each term for total phenolic content.	59
Table 4.18 Sum of squares and the percent contribution for each term for total flavonoid content.	59
Table 4.19 Sum of squares and the percent contribution for each term for antioxidant activity.	60
Table 4.20 Condition for factors optimizing polyphenol extraction.	61
Table 4.21 Experimental design and response for MAE optimization.	63
Table 4.22 Condition for factors in optimizing polyphenol extraction.	67
Table 4.23 Comparison between predicted and experimental value for optimum condition.	68
Table 5.1 Physical properties of spray dry powder with different encapsulating agent.	71

## LIST OF FIGURES

Figure 2.1 Scanning electron micrographs (100X) of Epimedium leaf sample. (A) Untreated leaf; (B) After ultrasonic assisted extraction by Zhang et al. (2009).	10
Figure 2.2 Transmission electron micrographs (3000×) of Epimedium leaf samples from (A), untreated leaf; (B), after ultrasonic-assisted extraction by Zhang et al., (2009).	11
Figure 2.3 Light micrographs of Epimedium leaf samples: (A) untreated leaf sample; (B) leaf sample after microwave irradiation by Zhang et al., (2011).	14
Figure 2.4 Active components in Phyllanthus Niruri extract.	15
Figure 2.5 Morphology of different type of microcapsules.	19
Figure 2.6 HPLC chromatogram of Phyllanthus Niruri extract from various authors.	23
Figure 3.1 Experimental work flow.	25
Figure 3.2 Phyllanthus Niruri Plant.	26
Figure 3.3 Spray Dryer and Schematic diagram of spray dryer.	32
Figure 3.4 Standard Calibration Curve of Gallic acid.	33
Figure 3.5 Standard Calibration Curve of Quercetin.	34
Figure 3.6 Standard Calibration Curve of Antioxidant Activity.	35
Figure 3.7 Standard Calibration curve of Phyllanthin.	36
Figure 3.8 Standard Calibration curve of Gallic Acid.	37
Figure 3.9 Standard Calibration curve of Quercetin.	37
Figure 4.1 UPLC chromatogram of P. niruri extract and chemical structures of the markers.	40
Figure 4.2 Identification of phyllanthin by matching UV spectra of sample to standard in Empower software library.	41
Figure 4.3 Identification of gallic acid by matching UV spectra of sample to standard in Empower software library.	41
Figure 4.4 Identification of quercetin by matching UV spectra of sample to standard in Empower software library.	41
Figure 4.5 Correlation of range of variables and predicted optimum condition for simultaneous maximum polyphenols recovery using the CCD model.	51
Figure 4.6 Predicted optimum condition for simultaneous maximum polyphenols recovery using CCD mode.	52
Figure 4.7 Correlation of range of variables and predicted optimum condition for simultaneous maximum polyphenols recovery using the CCD model.	65

Figure 4.8 Predicted optimum for simultaneous maximum polyphenols recovery using CCD model.	66
Figure 5.1 Particle size distribution at different formulation.	72
Figure 5.2 Surface morphology of the non-encapsulated and encapsulated spray dried powder. (A)-Without encapsulating agent; (B)- 10% MD; (C) 10% WPI; (D)- 10% WPI (1): MD (9).	74
Figure 5.3 Retention of phyllanthin, gallic acid and quercetin using different encapsulation strategies.	75

## LIST OF SYMBOLS

%	Percentage
°C	Degree Celsius
hr	Hour
$\lambda$	Wavelength
$\epsilon'$	Dielectric constant
kHz	KiloHertz
wt	Weight
$\tan\delta$	Dissipator factor
cm	Centimeter
mm	Milimeter
$\mu\text{m}$	Micrometer
kg	Kilogram
g	gram
ml	Mililiter



## LIST OF ABBREVIATIONS

AA	Antioxidant activity
ANOVA	Analysis of variance
BHA	Butylated hydroxyanisole
CCD	Central composite design
DE	Dextrose equivalent
DoE	Design of experiment
DPPH	2,2-diphenyl-1-picrylhydrazyl
DW	Dry weight
EtOH	Ethanol
Eup	Eupatorin
FESEM	Field emission scanning electron microscopy
GA	Gallic acid
H <sub>2</sub> O	Water
HPLC	High performance liquid chromatography
i.d.	Internal diameter
MAE	Microwave assisted extraction
MD	Maltodextrin

Min	Minute
PHY	Phyllanthin
QUE	Quercetin
RSM	Response surface methodology
Sec	Second
TFC	Total flavonoid content
TPC	Total phenolic content
UAE	Ultrasonic assisted extraction
UPLC	Ultra performances liquid chromatography
WPI	Whey protein isolate

## REFERENCES

- AbouZid, S. F., and Eklsherbeiny, G. M. 2008. Increase in flavonoids content in red onion peel by mechanical shredding. *Journal of Medical plant Research*. 82, 258-260.
- Akowuah, G. A., Ismail, Z., Norhayati, I. and Sadikun, A. 2005. Effect of extraction temperature on stability if major polyphenols and antioxidant activity of orthosiphon stamineus leaf. *Journal of herbs, spices and medical plants*. 16, 3-4.
- Al-dhabi, N. A., Ponmurugan, K., Maran, P. 2017. Development and validation of ultrasonic-assisted solid0liquid extraction on phenolic compounds from waste spent coffee grounds. *Ultrason. Sonochem*. 34, 206-213.
- Anderson, M. J., Whitcomb, P. J., Kraber, S. L., and Adams, W. 2009. Stat-Ease Handbook for Experimenters. Minneapolis: Stat-Ease, Inc.
- Anandharamkrishnan, C., Rielly, C. D. and Stapleym A. G. F. 2008. Loss of solubility of alpha-lactalbumin and beta-lactoglobulin during the spray drying of whey proteins. *LWT-Food Science and Technology*. 41(2), 270-277.
- Annamalai, A. and Lakshimi, P. T. V. 2009. HPTLC and HPLC Analysis of Bioactive Phyllanthin from Different Organs of Phyllanthus amarus. *Asian Journal of Biotechnology*. 1(4), 154-162.
- Araruna, S. M. 2008. Desenvolvimento, Padronização (CLAEDAD) e Avaliação Pré-Clinica do Extrato Seco por spray dried de Amburana cearensis A. C. Smith (Cumaru). Fortaleza. Dissertação de Mestrado, Programa de Pósgraduaçãoem Ciências Farmacêuticas, UFC.
- Baranauskiene, R., Venskutonis, P. R., Dewettinck, K., and Verhe, R. 2006. Properties of oregano (*Origanum vulgare* L.), citraonella (*Cymbopogon nardus* G.) and marjoram (*Majorana hortensis* L.) flavors encapsulated into milk protein-based matrices. *Food Research International*. 38, 989-994.
- Barbara, S., Dybowksi, F. and Krauze-Baranowska, M. 2015. Analysis of Securinega-type alkaloids from *Phyllanthus glaucus*. *Phytochemistry letter*. 11, 353-357.
- Barros, M.E., Lima, R., Mercuri, L. P., Matos, J. R., Schor, N., Boim, M. A. 2006. Effect of extract of *Phyllanthus niruri* on crystal deposition in experimental urolithiasis. *Urol Res*. 34, 351-357.
- Bhope, S. G., Kuber, V. V, Nagore, D. H., Gaikwad, P. S., and Patil M. J. 2013. Development and validation of RP-HPLC method for simultaneous analysis of andrograpolid, Phyllanthin, and Hypophyllanthin from herbal hepatoprotective formulation. *Acta Chromatographica*. 25(1), 159-169.
- Bylaite, E., Nylander, T., Venskutonis, R., and Jonsson, B. 2011. Emulsification of caraway essential oil in water by lecithin and beta-lactoglobulin: emulsion stability and properties of the formed oil- aqueous interface. *Colloids and Surface B: Biointerfaces*, 20(4), 327-340.

- Carrera, C., Ruiz-rodriguez. A., Palma, M., Barroso, C. G. 2012. Ultrasound assisted extraction of phenolic compounds from grapes. *Anal. Chim. Acta.* 723, 100-104.
- Chang, C., Yang, M., Wen, H. and Chern, J. 2002. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J. Food Drug Analysis.* 10, 178-182.
- Chatterjee, M., Sarkar, K., Sil, P.C. 2006. The protein isolate of the herb, *Phyllanthus niruri*, protects liver from nimesulide induced oxidative stress. *Pathophysiology* 13, 95–102.
- Chemat, F., Abert-Vian, M., Zill-e-Huma, Y-J. 2009. Microwave assisted separations: green chemistry in action. In: Perlman JT(ed) Green chemistry research trends. Nova Science Publisher, New York. pp 33-62.
- Chen, L., Song, D., Tian, Y., Ding, L., Yu, A. and Zhang, H. 2008. Application of on-line microwave sample-preparation techniques. *Trends Anal Chem.* 27, 151-159.
- Chen, M., Zhao, Y., and Yu, S. 2015. Optimisation of ultrasonic-assisted extraction of phenolic compounds, antioxidants, and anthocyanins from *sugar beet* molasses. *Food Chemistry.* 172, 543-550.
- Choi, K. O., Ryu, J., Kwak, H. S., and Ko, S. 2010. Spray-dried conjugated linoleic acid encapsulated with Maillard reaction products of whey proteins and maltodextrin. *Food Science and Biotechnology.* 19(4), 957-965.
- Cuoto, A. G., Kunzler, M, Spaniol, B., Magalhaes, P., Ortega, G. Petrovick, P. 2013. Chemical and technological evaluation of the *Phyllanthus niruri* aerial parts as a function of cultivation and harvesting conditions. *Brazilian Journal of Pharmacognosy.* 23(1), 36-43.
- Diezak, J. D. 1988. Microencapsulation and encapsulated ingredients. *Food Technology,* 42(4), 136-151.
- Dong, W., Yu, S., Deng, Y. and Pan, T. 2016. Screening of lignin patterns in *Schisandra* species using ultrasonic assisted temperature switch ionic liquid microextraction followed by UPLC-MS/MS analysis. *Journal of Chromatography B.* 1008, 45-49.
- El-Mahmood, A. M., Ogbonna, O. B. and Raji, M. 2010. The antibacterial activity of *Azadirachta indica* (Neem) seeds extracts against bacterial pathogens associated with eye and ear infections. *Journal of Medicinal Plants Research.* 4(14), 1414-1421.
- Eskilsson, C. S. and Bjorklund, E. 2000. Analytical-scale microwave-assisted extraction. *J Chromatogr A.* 902, 227-250.
- Fang, X., Wang, J., Wang, Y., Li, X., Zhou, H., and Zhu, L. 2014. Optimization of ultrasonic-assisted extraction of wedelolactone and antioxidant polyphenols from *Eclipta prostrata* L using response surface methodology. *Separation and Purification Technology.* 138, 55-64.

- Fang, Z., Zhang, M., Sun, Y. and Sun, J. 2007. Polyphenol oxidase from bayberry (*Myrica rubra* Sieb. et Zucc.) and its role in anthocyanin degradation. *Food Chemistry*. 103, 268-273.
- Falleh, H., Ksouri, R., Lucchessi, M., Abdelly, C., and Magne, C. 2012. Ultrasound-Assisted Extraction: Effect of Extraction Time and Solvent Power on the Levels of Polyphenols and Antioxidant Activity of *Mesembryanthemum edule* L. *Aizoaceae* Shoots. *Tropical Journal of Pharmaceutical Research*. 11(2), 243-249.
- Franco, J., Sineiro, M., Rubilar, M., Sanchez, M., Jerez, M., Pinelo, M. and Nunez, M. J. 2008. Polyphenols from plant materials: extraction and antioxidant power. *Electronic Journal of Environment, Agriculture and Food Chemistry*. 7, 3210-3216.
- Fernandes, L. P., Candido, R. C., and Oliveira, W. P. 2012. Spray drying microencapsulation of Lippia sidoides extracts in carbohydrate blends. *Food and Bioproducts Processing*. 90(3), 425-432.
- Garcia-castello, E. M., Rodriguez-lopez, A. D., Mayor, L., Ballesteros, R., Conidi, C., Cassano, A. 2015. Optimization of conventional and ultrasound assisted extraction of flavonoids from grapefruit (*Citrus paradisi* L.) solid wastes. *LWT Food Sci. Technol*. 64, 1114–1122.
- Gibbs, B. F., Kermasha, S., Alli, I., and Mulligan, C. N. 1999. Encapsulation in the food industry A review. *International Journal of Food Science and Nutrition*. 50, 213-224.
- Ghosal. S., Veeraragavan, M., and Kalidindi, S. R. (2013). *U.S. Patent No. 20130122119A1*. New Brunswick. DC: U.S Patent and Office Trademark Office.
- Gouin, S. 2004. Micro-encapsulation: Industrial appraisal of existing technologies and trends. *Trends in Food Science and Technology*. 15, 330-347.
- Harish, R., Shivanandappa, T. 2006. Antioxidant activity and hepatoprotective potential of *Phyllanthus niruri* . *Food Chem*. 95, 180–185.
- Hashemi, A. M. B., Khaneghah, A. M., and Akbarirad, H. 2016. The effect of Amplitudes Ultrasound-Assisted Solvent Extraction and Pretreatment Time on the Yield and Quality of Pistacia Khiniuk Hull Oil. *Journal of Oleo Science*. 65, 733-738.
- Ichoo, M., Chirdchupunseree, H., Pramyothin, P., and Jianmongkol, S. 2011. Endothelium-independent effects of phyllanthin and hypophyllanthin on vascular tension. *Fitoterapia*. 82, 1231-1236.
- Islam, A., Selvan, T., Mazumder, U. K., Gupta, M., and Ghosal, S. 2008. Antitumour effect of phyllanthin and hypophyllanthin from phyllanthus amarus against ehrlich ascites carcinoma in mice. *Pharmacologyonline*. 2, 796-807.
- Jaiswal, A.K. and Abu-Ghannam, N. 2013. Degradation kinetic modelling of color, texture, polyphenols and antioxidant capacity of York cabbage after microwave processing. *Food Research International*. 53, 125-133.

- Jassie L, Reveszm R., Kierstead T., Hasty, E., Metz, S. 1997. In: Kingston HM, Haswell SJ (eds.). Microwave-enhanced chemistry. American Chemical Society, Washington, DC. Pp569.
- Karimi, E., Jaafar, H.Z.E., Aziz, M.A., Taheri, S. and AzadiGonbad, R. 2014. Genetic relationship among *Labisia pumila* (Myrsinaceae) species based on ISSR-PCR. *Genetics and Molecular Research*. 13(2): 3301-3309. doi: <http://dx.doi.org/10.4238/2014.April.29.8>
- Kassuya, C. A. L., Silverstre, A., Menezes-de-Lima, O. J., Marotta, D. M., Lucia, V. and Calixto, J. 2006. Antiinflammatory and antiallodynic actions of the lignin niranthin isolated from *Phyllanthus amarus*. *European Journal of Pharmacology*. 546: 182-188.
- Khajeh M, Moghaddam, A.R. and Sanchooli, E. 2009. Application of Doehlert design in the optimization of microwave assisted extraction for determination of zinc and copper in cereal samples using FAAS. *Food Anal Methods*. 3(3): 133-137.
- Krishnaiah, D., Sarbatly, R. and Nithyanandam, R. 2012. Microencapsulation of Morinda citrifolia L. extract by spray drying. *Chemical Engineering Research and Design*. 90(5): 622-632
- King, A. H. 1995. Encapsulation of food ingredients: A review of available technology, focusing on hydrocolloids. In S. J. Risch, & G.A. Reineccius (Eds.), Encapsulation and controlled release of food ingredients. ACS symposium series (Vol. 590, pp. 26-39). Washington, DC: American Chemical Society.
- Li, J., Zu, Y. G, Fu, Y. J, Yang, Y. C, Li, S. M., Li, Z. N., and Wink. M. 2010. Optimization of microwave-assisted extraction of triterpene saponins from defatted residue of yellow horn (*Xanthoceras sorbifolia* Bunge.) kernel and evaluation of its antioxidant activity. *Innov food Sci Emerg Technol*. 11: 637-664.
- Li, Y., Li, S., Lin, S., Zhang, J., Zhao, C. and Li, H. 2017. Microwave-Assisted Extraction of Natural Antioxidants from the Exotic *Gordonia axillaris* Fruit: Optimization and Identification of Phenolic compounds. *Molecule*. 22(9); 1481.
- Maity, S., Chatterjee, S., Variyar, P. S., Sharma, A., Adhikari, S., Mazumder, S. 2013. Evaluation of antioxidant activity and characterization of phenolic constituents of *Phyllanthus amarus* root. *J. Agric. Food Chem*. 61: 3443-3450.
- Mandal, V, Mohan, Y. and Hemalath, S. 2007. Microwave assisted extraction-an innovative and promising extraction tool for medical plant research. *Phcog Rev*. 1(1): 7-18.
- Markom, M., Hasan, M., Daud, W. R. W., Singh, H., and Jahim, H. M. (2007). Extraction of hydrolysable tannins from *Phyllanthus niruri* Linn.: Effects of solvents and extraction methods. *Separation and Purification Technology*. 52: 487-496.
- Martendal, E., Budziak, D., and Carasek, E. 2007. Application of fractional factorial experimental and Box-Behnken designs for optimization of single-drop microextraction of 2,4,6-trichloroanisole and 2,4,6-tribromoanisole from wine

- samples. *Journal of Chromatography A*, 1148: 131–136.
- Montgomery, D. C., and Runger, G. C. 2010. Applied Statistics and Probability for Engineers. John Wiley & Sons
- Mediani, A., Abas, F., Tan, C. P., Ismail, I. S., Shaari, K., Ismail, A., and Lajis, N. H. 2015. Phytochemical and biological features of *Phyllanthus Niruri* and *Phyllanthus urinaria* harvested at different growth stages revealed by H NMR-based metabolomics. *Industrial Crops and Products*. 77: 602-613.
- Mee, R. (2009). A comprehensive guide to factorial two-level experimentation. London: Springer.
- Metrouh-Amir, H., Durte, C. M. and Maiza, F. 2015. Solvent effect on total phenolic contents, antioxidant, and antibacterial activities of *Matricaria pubescens*. *Industrial Crops and Product*. 67; 249-256.
- Meullemiestre, A., Petitcolas, E., Maache-rezzoug, Z., Chemat, F. Rezzoug, S. A. 2016. Impact of ultrasound on solid-liquid extraction of phenolic compounds from maritime pine sawdust waste. Kinetics, optimization and large scale experiments. *Ultrason. Sonochem*. 28, 230-239.
- Ministry of Health Malaysia. (2015). Health Facts. Retrieved from [http://vlib.moh.gov.my/cms/documentstorage/com.tms.cms.document.Document\\_t\\_2e8efe25-a0188549-d5315d00-e9ca9560/KKM\\_HEALTH\\_FACTS\\_2015.pdf](http://vlib.moh.gov.my/cms/documentstorage/com.tms.cms.document.Document_t_2e8efe25-a0188549-d5315d00-e9ca9560/KKM_HEALTH_FACTS_2015.pdf)
- Miranda, M., Vega-Galvez, A., Lopez, J., Parada, G., Sanders, M., Aranda, M., Uribe, E., Di Scale, K. 2010. Impact of air drying temperature on nutritional properties, total phenolic content and antioxidant capacity of quinoa seeds (*Chenopodium quinoa* Wild.), *Industrial Crops and Products*. 32: 258-263.
- Moreira, J., Klein-Junior, L. C., Filho, V. C., Buzzi, F. C. 2013. Anti-hyperalgesic activity of corilagin, a tannin isolated from *Phyllanthus niruri* L. (Euphorbiaceae). *Journal of Ethnopharmacology*. 146, 318-323.
- Murugaiyah, V., and Chan, K, L. 2007. Analysis of lignans from *Phyllanthus niruri* L. in plasma using a simple HPLC method with fluorescence detection and its application in a pharmacokinetic study. *Journal of Chromatography*. 852, 138-144.
- Nikam, P. S., Nikam, S. V., Sontakke, A.V., Khanwelkar, C. C. 2011. Role of *Phyllanthus amarus* treatment in hepatitis-C . *Biomed. Res*. 22: 319-322.
- Nishiura, J.L., Campos, A.H., Boim, M.A., Heilberg, I.P., Schor, N. 2004. *Phyllanthus niruri* normalizes elevated urinary calcium levels in calcium stone forming (CSF) patients. *Urol*. 32, 362–366.
- Nguang, S. L, Yeong Y. L, Pang, S. F., & Gim bun. J. 2016. Ultrasonic As-sisted Extraction on Phenolic and Flavonoid Content from *Phyllanthus niruri* Plant. *Indian Journal of Science and Technology* 10(2), 1-5.

- Noor, N.M., Annie, G., Hussain, N.H.N., Kadir, A.A., Hassan, I.I., Sooi, L.K., Ismail, S.B., Mohamed, W.H.W., Ekehard, L. and Abas, A. 2014. Efficacy and safety of *Labisia pumila* var *alata* water extract among pre and postmenopausal women. *Journal of Medicinal Food*. 17(8): 929938. doi:10.1089/jmf.2013.2953.
- Notka, F., Meier, G. R., Wagner, R. 2004. Concerted inhibitory activities of *Phyllanthus amarus* on HIV replication *in vitro* and *ex vivo*. *J. Antivir. Res.* 64, 93-102.
- Ncube, N. S., Afolayan, A. J. and Okoh, A. I. 2008. Assessment techniques of antimicrobial properties of natural compounds of plant origin: current methods and future trends. *African Journal of Biotechnology*. 7 :1797-1806.
- Pan, X., Niu, G. and Liu, H. 2003. Microwave assisted extraction of tea polyphenols and tea caffeine from green tea leaves. *Chem Eng Process*. 42: 129-133.
- Pang, S.F., Lau, M.J., Gimbun, J., Yusoff, M.M. 2012. Extraction of flavonoids, antioxidant and phenolic content from *O. stamineus*, The National Conference on Postgraduate Research.
- Patel, J. R., Tripathi, P., Sharma, V., Chauhan, N. S., Dixit, V. K. 2011. *Phyllanthus amarus*: ethanomedicinal uses, phytochemistry and pharmacology: a review. *J. Ethnopharmacol.* 138: 286-313.
- Poh-Hwa, T., Yoke-Kqueen, C, Indu Bala, J. and Son, R. 2011. Bioprotective properties of three Malaysia *Phyllanthus* species: An Investigation of the Antioxidant and Antimicrobial Activities. *International Food Research Journal*. 18, 887-893.
- Quy, D. D., Angkawijaya, A. E., and Tran-Nguyen, P. L. 2014. Effect of extraction solvent and total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatic*. *Journal of Food and Drug Analysis*. 296-301
- Raphal, K.R., Kuttan, R. 2003. Inhibition of experimental gastric lesion and inflammation by *Phyllanthus amarus* extract. *J. Ethnopharmacol.* 87, 193–197.
- Reineccius, G. A. 1988. Spray drying of food flavours. In G. A. Reineccius, & S. J. Risch (Eds.), *Flavour encapsulation* (pp. 55-66). Washington, DC: American Chemical Society, 55-66.
- Rosenberg, M. and Sheu, T. Y. 1998. Microencapsulation of volatiles by spray-drying in whey protein based wall systems. *International Dairy Journal*. 6, 273-284.
- Routray, W. and Orsat, W. 2011. Microwave-assisted extraction of flavonoids: a review. *Food Bioprocess Technol.* 5, 1-16.
- Sarkar, K., Ghosh, A., Sil, P.C. 2005. Preventive and curative role of a 43kD protein from the leaves of the herb *Cajanus indicus* L on thioacetamide-induced hepatotoxicity *in vivo*. *Hepatology*. 33, 39–49
- Sarkar, M.K., Sil, P.C. 2010. Prevention of tertiary butyl hydroperoxide induced oxidative impairment and cell death by a novel antioxidant protein molecule isolated from the herb, *Phyllanthus niruri*. *Toxicol. In Vitro* 24, 1711–1719.



- Shahidi, F., and Han, X. Q. 1993. Encapsulation of food ingredients. *Critical Review in Food Science and Nutrition*. 33, 501-547.
- Sharma, P., Parmar, J., Verma, P., Sharma, P., & Goyal, P. K. 2009. Anti-tumor activity of *Phyllanthus niruri* (a medical plant) on chemical-induced skin carcinogenesis in mice. *Asian Pac J Cancer Prec* 10(6), pp 1089-1094.
- Sherwin, E. R. 1978. Oxidation and antioxidants in fat and oil processing. *Journal of the American Oil Society*. 55, 809-814.
- Soelaiman, I.N., Das, S., Shuid, A.N., Mo, H. and Mohamed, N. 2013. Use of medicinal plants and natural products for treatment of osteoporosis and its complications. *Evidence-Based Complementary and Alternative Medicine*. 2013: 764701. doi: <http://dx.doi.org/10.1155/2013/764701>
- Sólyom, K., Solá, R., Cocero, M.J., Mato, R.B. 2014. Thermal degradation of grape marc polyphenols. *Food Chemistry*. 159, 361-366.
- Song, J., Li, D., Liu, C., and Zhang, Y. 2011. Optimized microwave-assisted activity. *Inno Food Sci Emerg Technol* 12, 282-287.
- Soottitantawat, A., Takayama, K., Okamura, K., Muranaka, D., Yoshii, H. and Furutani T. 2005b. Microencapsulation of L-menthol by spray drying and its release characteristics. *Innovative Food Science and Emerging Technologies*. 6, 163-170.
- Sousa, A. D., Maia, A. I. V., Rodrigues, T. H.S., Canuto, K. M., Ribeiro, P. R. V., Pereira, R. C. A., Viera, R. F. and Brito, E. S. 2016. Ultrasonic-assisted and pressurized liquid extraction of phenolic compounds from *Phyllanthus amarus* and its composition evaluation by UPLC-QTOF. *Industrial Crops and Products*. 79, 91-103.
- Swartz, M. E. 2005. UPLC An introduction and review. *J. Liq. Chromatogr*. 28 1253-1263.
- Tang, Y. Q., and Shamala, D. S. 2011. Evaluation of *Phyllanthus*, for Its Anti-Cancer Properties, Prostate Cancer - From Bench to Bedside, Dr. Philippe E. Spiess (Ed.), ISBN: 978-953-307-331-6, InTech, Available from: <http://www.intechopen.com/books/prostate-cancer-from-bench-to-bedside/evaluation-ofphyllanthus-for-its-anti-cancer-properties>
- Tan, H. T., Lee, K. T., and Mohamed, A. R. 2011. Pretreatment of lignocellulosic palm biomass using a solvent-ionic liquid [BMIM]Cl for glucose recovery: An optimisation study using response surface methodology. *Carbohydrate Polymers*, 83, 1862–1868.
- Talebi, M., Ghassempour, A., Talebpour, Z., Rassouli, A. and Dolatyari, L. 2004. Optimization of the extraction of paclitaxel from *Taxus baccata* L. by the use of microwave energy. *J Sep Sci*. 27, 1130-1136.
- Tara, S., Arul. A., Amita, S. and Sudhakar. 2010. Effect of *Phyllanthus Niruri*. Linn on burn wound in rats. *Asian Pacific Journal of Tropical Medicine*. 105-108

- Teixeira, M. I., Andrade, J. R., Farina, M. and Rocha-Leao, M. H. M. 2004. Characterization of short chain fatty acid microcapsules produced by spray drying. *Materials Science and Engineering*. 24, 653-658.
- Thostenson, E. T. and Chou, T. W. 1999. Microwave processing: fundamentals and applications. *Compos Part A Appl S*. 30(9),1055-1071.
- Trabelsi, N., Megdiche. M., Ksouri, R., Falleh, H., Oueslati, S., SoumayaB., Hajlaoui, H. and Abdelly, C. 2010. Solvent effects on phenolic content and biological activities of the halophyte *Limoniastrum monopetalum* leaves. *LWT-Food Science and Technology*. 43, 632-639.
- Verbeyst, L., Oey, I., Van der Planchen, I., Hendrickx, M., Loey, A. V., 2010. Kinetic study on the thermal and pressure degradation of anthocyanins in strawberries. *Food Chemistry*. 123, 269-274.
- Wang, L. and Weller, C. L. 2006. Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science & Technology*. 17, 300-312.
- Wang, X., Wu, Y., Chen, G., Yue, W., Liang, Q., Wu, Q. 2013. Optimization of ultrasonic assisted extraction of phenolic compounds from Sparganji rhizome with response surface methodology. *Ultrasonic. Sonoch*. 20, 846-854.
- Watanabe, Y., Fang, X., Adachi, S., Fukami, H., and Matsuno, R. 2004. Oxidation of 6-O-arachidonoyl-ascorbate microencapsulated with a polysaccharide by spray-drying. *Lebensmittel-Wissenschaft und-Technologie*. 37: 395-400.
- Xi, J. 2009. Caffeine extraction from green tea leaves assisted by high pressure processing. *Journal of Food Engineering*. 94, 105-109.
- Xia, Y, Luo, H, Liu, J. P., & Gluud, C. 2011. Phyllanthus species for chronic hepatitis B virus infection *Cochrane Database Syst Rev* 13(4) doi: 10.1002/14651858.CD008960.pub2.
- Xie, Y. L., Zhou, H. M., Liang, X. H, He, B. S. and Han, X. X. 2010. Study on the Morphology, Particle Size and Thermal Properties of Vitamin A Microencapsulated by Starch Octenylsuccinate. *Agricultural Science in China*. 9(7): 1058-1060.
- Young, S. L., Sarda, X. and Rosenberg, M. 1993. Microencapsulating Properties of Whey Protein. 2. Combination of Whey Protein with carbohydrate. *Journal of Dairy Science*. 76(10): 2878-2885.
- Zhang, H. F., Yang, X. H., Zhao, L. D., and Wang, Y. 2009. Ultrasonic-assisted extraction of *Epimedium* and extraction mechanism. *Innovative Food Science and Engineering Technologies*. 10: 54-60.
- Zhang, H. F., Yang, X. H. and Wang, Y. 2011. Microwave assisted extraction of secondary metabolites from plants: Current status and future directions. *Trends in Food Science & Technology*. 22(12): 672-688.

Zhi-feng, F., Zong-cai, T., Lu, Z., Hui, W., Qing-hui, W. and Tao, H. 2016. Antioxidant activities and polyphenols of sweet potato (*Ipomoea batatas* L.) leaves extracted with solvents of various polarities. *Food Bioscience*. 15, 11-18

Zlotorzynski, A. 1995. The application of microwave radiation to analytical and environmental chemistry. *Crit Rec Anal Chem*. 25: 43-7